



VICTORIA UNIVERSITY
MELBOURNE AUSTRALIA

Inequalities for α -fractional differentiable functions

This is the Published version of the following publication

Chu, Y-M, Adil Khan, M, Ali, T and Dragomir, Sever S (2017) Inequalities for α -fractional differentiable functions. Journal of Inequalities and Applications. ISSN 1025-5834

The publisher's official version can be found at
<https://journalofinequalitiesandapplications.springeropen.com/articles/10.1186/s13660-017-1371-6>

Note that access to this version may require subscription.

Downloaded from VU Research Repository <https://vuir.vu.edu.au/35896/>

RESEARCH

Open Access



Inequalities for α -fractional differentiable functions

Yu-Ming Chu^{1*}, Muhammad Adil Khan², Tahir Ali² and Sever Silvestru Dragomir³

*Correspondence:
chuyuming2005@126.com
¹Department of Mathematics,
Huzhou University, Huzhou, 313000,
China
Full list of author information is
available at the end of the article

Abstract

In this article, we present an identity and several Hermite-Hadamard type inequalities for conformable fractional integrals. As applications, we establish some inequalities for certain special means of two positive real numbers and give the error estimations for the trapezoidal formula.

MSC: 26D15; 26A51; 26A33

Keywords: convex function; Hermite-Hadamard inequality; fractional derivative; fractional integral; special mean; trapezoidal formula

1 Introduction

A real-valued function $\psi : I \subseteq \mathbb{R} \rightarrow \mathbb{R}$ is said to be convex on I if the inequality

$$\psi(\theta\xi + (1-\theta)\zeta) \leq \theta\psi(\xi) + (1-\theta)\psi(\zeta) \quad (1.1)$$

holds for all $\xi, \zeta \in I$ and $\theta \in [0, 1]$. ψ is said to be concave on I if inequality (1.1) is reversed.

Let $\psi : I \subseteq \mathbb{R} \rightarrow \mathbb{R}$ be a convex function on the interval I , and $c_1, c_2 \in I$ with $c_1 < c_2$. Then the double inequality

$$\psi\left(\frac{c_1 + c_2}{2}\right) \leq \frac{1}{c_2 - c_1} \int_{c_1}^{c_2} \psi(\xi) d\xi \leq \frac{\psi(c_1) + \psi(c_2)}{2} \quad (1.2)$$

is known in the literature as the Hermite-Hadamard inequality for convex functions [1–3]. Both inequalities hold in the reversed direction if ψ is concave on the interval I . In particular, many classical inequalities for means can be derived from (1.2) for appropriate particular selections of the function ψ .

Recently, the improvements, generalizations, refinements and applications for the Hermite-Hadamard inequality have attracted the attention of many researchers [4–22].

Dragomir and Agarwal [23] proved the following results connected with the right hand part of (1.2).

Theorem 1.1 (See [23], Lemma 2.1) *Let $\psi : I^\circ \subseteq \mathbb{R} \rightarrow \mathbb{R}$ be a differentiable mapping on I° . Then the identity*

$$\frac{\psi(c_1) + \psi(c_2)}{2} - \frac{1}{c_2 - c_1} \int_{c_1}^{c_2} \psi(\xi) d\xi = \frac{c_2 - c_1}{2} \int_0^1 (1 - 2\theta) \psi'(\theta c_1 + (1 - \theta)c_2) d\theta$$

holds for all $c_1, c_2 \in I^\circ$ with $c_1 < c_2$ if $\psi' \in L[c_1, c_2]$, where I° denotes the interior of I .

Theorem 1.2 (See [23], Theorem 2.2) *Let $\psi : I^\circ \subseteq \mathbb{R} \rightarrow \mathbb{R}$ be a differentiable mapping on I° . Then the inequality*

$$\left| \frac{\psi(c_1) + \psi(c_2)}{2} - \frac{1}{c_2 - c_1} \int_{c_1}^{c_2} \psi(\xi) d\xi \right| \leq \frac{(c_2 - c_1)(|\psi'(c_1)| + |\psi'(c_2)|)}{8}$$

holds for $c_1, c_2 \in I^\circ$ with $c_1 < c_2$ if $|\psi'|$ is convex on $[c_1, c_2]$.

Making use of Theorem 1.1, Pearce and Pečarić [24] established Theorem 1.3 as follows.

Theorem 1.3 (See [24], Theorem 1) *Let $c_1, c_2 \in I \subseteq \mathbb{R}$ with $c_1 < c_2$, $\psi : I^\circ \subseteq \mathbb{R} \rightarrow \mathbb{R}$ be a differentiable mapping on I° and $q \geq 1$. Then the inequality*

$$\left| \frac{\psi(c_1) + \psi(c_2)}{2} - \frac{1}{c_2 - c_1} \int_{c_1}^{c_2} \psi(\xi) d\xi \right| \leq \frac{c_2 - c_1}{4} \left[\frac{|\psi'(c_1)|^q + |\psi'(c_2)|^q}{2} \right]^{1/q}$$

is valid if the mapping $|\psi'|^q$ is convex on the interval $[c_1, c_2]$.

Next, we recall several elementary definitions and important results in the theory of conformable fractional calculus, which will be used throughout the article, we refer the interested reader to [25–32].

The conformable fractional derivative of order $0 < \alpha \leq 1$ for a function $\psi : (0, \infty) \rightarrow \mathbb{R}$ at $\xi > 0$ is defined by

$$D_\alpha(\psi)(\xi) = \lim_{\epsilon \rightarrow 0} \frac{\psi(\xi + \epsilon \xi^{1-\alpha}) - \psi(\xi)}{\epsilon},$$

and the fractional derivative at 0 is defined as $D_\alpha(\psi)(0) = \lim_{\xi \rightarrow 0^+} D_\alpha(\psi)(\xi)$.

The (left) fractional derivative starting from c_1 of a function $\psi : [c_1, \infty) \rightarrow \mathbb{R}$ of order $0 < \alpha \leq 1$ is defined by

$$D_\alpha^{c_1}(\psi)(\xi) = \lim_{\epsilon \rightarrow 0} \frac{\psi(\xi + \epsilon(\xi - c_1)^{1-\alpha}) - \psi(\xi)}{\epsilon},$$

and we write $D_\alpha^{c_1}(\psi) = D_\alpha^0(\psi) = D_\alpha(\psi)$ if $c_1 = 0$. For more details see [26].

Let $\alpha \in (0, 1]$ and ψ, ϕ be α -differentiable at $\xi > 0$. Then we have

$$\frac{d_\alpha}{d_\alpha \xi}(\xi^n) = n \xi^{n-\alpha}, \quad n \in \mathbb{R},$$

$$\frac{d_\alpha}{d_\alpha \xi}(c) = 0, \quad c \in \mathbb{R},$$

$$\frac{d_\alpha}{d_\alpha \xi}(c_1 \psi(\xi) + c_2 \phi(\xi)) = c_1 \frac{d_\alpha}{d_\alpha \xi}(\psi(\xi)) + c_2 \frac{d_\alpha}{d_\alpha \xi}(\phi(\xi)), \quad c_1, c_2 \in \mathbb{R},$$

$$\frac{d_\alpha}{d_\alpha \xi}(\psi(\xi) \phi(\xi)) = \psi(\xi) \frac{d_\alpha}{d_\alpha \xi}(\phi(\xi)) + \phi(\xi) \frac{d_\alpha}{d_\alpha \xi}(\psi(\xi)),$$

$$\begin{aligned}\frac{d_{\alpha}}{d_{\alpha}\xi}\left(\frac{\psi(\xi)}{\phi(\xi)}\right) &= \frac{\phi(\xi)\frac{d_{\alpha}}{d_{\alpha}\xi}(\psi(\xi)) - \psi(\xi)\frac{d_{\alpha}}{d_{\alpha}\xi}(\phi(\xi))}{(\phi(\xi))^2}, \\ \frac{d_{\alpha}}{d_{\alpha}\xi}(\psi(\phi)(\xi)) &= \psi'(\phi(\xi))\frac{d_{\alpha}}{d_{\alpha}\xi}(\phi(\xi)),\end{aligned}\quad (1.3)$$

where ψ is differentiable at $\phi(\xi)$ in equation (1.3). In particular,

$$\frac{d_{\alpha}}{d_{\alpha}\xi}(\psi(\xi)) = \xi^{1-\alpha} \frac{d}{d\xi}(\psi(\xi))$$

if ψ is differentiable.

Let $\alpha \in (0, 1]$ and $0 \leq c_1 < c_2$. A function $\psi : [c_1, c_2] \rightarrow \mathbb{R}$ is said to be α -fractional integrable on $[c_1, c_2]$ if the integral

$$\int_{c_1}^{c_2} \psi(\xi) d_{\alpha}\xi = \int_{c_1}^{c_2} \psi(\xi) \xi^{\alpha-1} d\xi$$

exists and is finite. All the α -fractional integrable functions on $[c_1, c_2]$ are denoted by $L_{\alpha}^1([c_1, c_2])$.

It is well known that

$$\int_{c_1}^{c_2} \psi(\xi) D_{\alpha}^{c_1}(\phi)(\xi) d_{\alpha}\xi = \psi\phi|_{c_1}^{c_2} - \int_{c_1}^{c_2} \phi(\xi) D_{\alpha}^{c_1}(\psi)(\xi) d_{\alpha}\xi$$

if $\psi, \phi : [c_1, c_2] \rightarrow \mathbb{R}$ are two functions such that $\psi\phi$ is differentiable.

Very recently, Anderson [33] established a Hermite-Hadamard type inequality for fractional differentiable functions as follows.

Theorem 1.4 *Let $\alpha \in (0, 1]$ and $\psi : [c_1, c_2] \rightarrow \mathbb{R}$ be an α -fractional differentiable function. Then the inequality*

$$\frac{\alpha}{c_2^{\alpha} - c_1^{\alpha}} \int_{c_1}^{c_2} \psi(\xi) d_{\alpha}\xi \leq \frac{\psi(c_1) + \psi(c_2)}{2} \quad (1.4)$$

holds if $D_{\alpha}(\psi)$ is increasing on $[c_1, c_2]$. Moreover, if the function ψ is decreasing on $[c_1, c_2]$, then one has

$$\psi\left(\frac{c_1 + c_2}{2}\right) \leq \frac{\alpha}{c_2^{\alpha} - c_1^{\alpha}} \int_{c_1}^{c_2} \psi(\xi) d_{\alpha}\xi. \quad (1.5)$$

Remark 1.5 We clearly see that inequalities (1.4) and (1.5) reduce to inequality (1.2) if $\alpha = 1$.

The main purpose of the article is to present an identity and several Hermite-Hadamard type inequalities for conformable fractional integrals, establish some inequalities for certain special means of two positive real numbers and give the error estimations for the trapezoidal formula.

2 Main results

In order to prove our main results we need a lemma, which we present in this section.

Lemma 2.1 *Let $\alpha \in (0, 1]$, $c_1, c_2 \in \mathbb{R}$ with $0 \leq c_1 < c_2$ and $\psi : [c_1, c_2] \rightarrow \mathbb{R}$ be an α -fractional differentiable function on (c_1, c_2) . Then the identity*

$$\begin{aligned} & \frac{\psi(c_1) + \psi(c_2)}{2} - \frac{\alpha}{c_2^\alpha - c_1^\alpha} \int_{c_1}^{c_2} \psi(\xi) d_\alpha \xi \\ &= \frac{(c_2 - c_1)}{2(c_2^\alpha - c_1^\alpha)} \left[\int_0^1 ((\theta c_1 + (1 - \theta)c_2)^{2\alpha-1} - c_2^\alpha (\theta c_1 + (1 - \theta)c_2)^{\alpha-1}) \right. \\ & \quad \times D_\alpha(\psi)(\theta c_1 + (1 - \theta)c_2) \theta^{1-\alpha} d_\alpha \theta \\ & \quad + \int_0^1 ((\theta c_1 + (1 - \theta)c_2)^{2\alpha-1} - c_1^\alpha (\theta c_1 + (1 - \theta)c_2)^{\alpha-1}) \\ & \quad \times D_\alpha(\psi)(\theta c_1 + (1 - \theta)c_2) \theta^{1-\alpha} d_\alpha \theta \left. \right] \end{aligned}$$

holds if $D_\alpha(\psi) \in L_\alpha^1([c_1, c_2])$.

Proof Let $\xi = \theta c_1 + (1 - \theta)c_2$. Then making use of integration by parts, we get

$$\begin{aligned} & \int_0^1 ((\theta c_1 + (1 - \theta)c_2)^{2\alpha-1} - c_2^\alpha (\theta c_1 + (1 - \theta)c_2)^{\alpha-1}) D_\alpha(\psi)(\theta c_1 + (1 - \theta)c_2) d\theta \\ & \quad + \int_0^1 ((\theta c_1 + (1 - \theta)c_2)^{2\alpha-1} - c_1^\alpha (\theta c_1 + (1 - \theta)c_2)^{\alpha-1}) D_\alpha(\psi)(\theta c_1 + (1 - \theta)c_2) d\theta \\ &= \int_0^1 ((\theta c_1 + (1 - \theta)c_2)^\alpha - c_2^\alpha) \psi'(\theta c_1 + (1 - \theta)c_2) d\theta \\ & \quad + \int_0^1 ((\theta c_1 + (1 - \theta)c_2)^\alpha - c_1^\alpha) \psi'(\theta c_1 + (1 - \theta)c_2) d\theta \\ &= ((\theta c_1 + (1 - \theta)c_2)^\alpha - c_2^\alpha) \frac{\psi(\theta c_1 + (1 - \theta)c_2)}{c_1 - c_2} \Big|_0^1 \\ & \quad - \int_0^1 \alpha (\theta c_1 + (1 - \theta)c_2)^{\alpha-1} (c_1 - c_2) \frac{\psi(\theta c_1 + (1 - \theta)c_2)}{c_1 - c_2} d\theta \\ & \quad + ((\theta c_1 + (1 - \theta)c_2)^\alpha - c_1^\alpha) \frac{\psi(\theta c_1 + (1 - \theta)c_2)}{c_1 - c_2} \Big|_0^1 \\ & \quad - \int_0^1 \alpha (\theta c_1 + (1 - \theta)c_2)^{\alpha-1} (c_1 - c_2) \frac{\psi(\theta c_1 + (1 - \theta)c_2)}{c_1 - c_2} d\theta \\ &= \frac{1}{c_2 - c_1} \left[(c_2^\alpha - c_1^\alpha) \psi(c_1) - \alpha \int_{c_1}^{c_2} \psi(\xi) d_\alpha \xi \right] \\ & \quad + \frac{1}{c_2 - c_1} \left[(c_2^\alpha - c_1^\alpha) \psi(c_2) - \alpha \int_{c_1}^{c_2} \psi(\xi) d_\alpha \xi \right] \\ &= \frac{c_2^\alpha - c_1^\alpha}{c_2 - c_1} (\psi(c_1) + \psi(c_2)) - \frac{2\alpha}{c_2 - c_1} \int_{c_1}^{c_2} \psi(\xi) d_\alpha \xi. \end{aligned} \tag{2.1}$$

Therefore, Lemma 2.1 follows easily from (2.1). \square

Remark 2.2 We clearly see that the identity given in Lemma 2.1 reduces to the identity given in Theorem 1.1 if $\alpha = 1$.

Theorem 2.3 Let $\alpha \in (0, 1]$, $c_1, c_2 \in \mathbb{R}$ with $0 \leq c_1 < c_2$ and $\psi : [c_1, c_2] \rightarrow \mathbb{R}$ be an α -differentiable function. Then the inequality

$$\left| \frac{\psi(c_1) + \psi(c_2)}{2} - \frac{\alpha}{c_2^\alpha - c_1^\alpha} \int_{c_1}^{c_2} \psi(\xi) d_\alpha \xi \right| \leq \frac{c_2 - c_1}{2(c_2^\alpha - c_1^\alpha)} \left[\frac{(|\psi'(c_1)| + |\psi'(c_2)|)(5c_2^\alpha - 7c_1^\alpha + c_1c_2^{\alpha-1} + c_1^{\alpha-1}c_2)}{12} \right] \quad (2.2)$$

holds if $D_\alpha(\psi) \in L_\alpha^1([c_1, c_2])$ and $|\psi'|$ is convex on $[c_1, c_2]$.

Proof It follows from Lemma 2.1 and the convexities of the functions $\xi \rightarrow \xi^{\alpha-1}$ and $\xi \rightarrow -\xi^\alpha$ on $(0, \infty)$ together with the convexity of $|\psi'|$ on $[c_1, c_2]$ that

$$\begin{aligned} & \left| \frac{\psi(c_1) + \psi(c_2)}{2} - \frac{\alpha}{c_2^\alpha - c_1^\alpha} \int_{c_1}^{c_2} \psi(\xi) d_\alpha \xi \right| \\ & \leq \frac{c_2 - c_1}{2(c_2^\alpha - c_1^\alpha)} \left[\int_0^1 ((\theta c_1 + (1 - \theta)c_2)^\alpha - c_1^\alpha) |\psi'(\theta c_1 + (1 - \theta)c_2)| d\theta \right. \\ & \quad \left. + \int_0^1 (c_2^\alpha - (\theta c_1 + (1 - \theta)c_2)^\alpha) |\psi'(\theta c_1 + (1 - \theta)c_2)| d\theta \right] \\ & \leq \frac{c_2 - c_1}{2(c_2^\alpha - c_1^\alpha)} \left[\int_0^1 ((\theta c_1 + (1 - \theta)c_2)^{\alpha-1} (\theta c_1 + (1 - \theta)c_2 - c_1^\alpha) |\psi'(\theta c_1 + (1 - \theta)c_2)| d\theta \right. \\ & \quad \left. + \int_0^1 (c_2^\alpha - ((1 - \theta)c_1^\alpha + \theta c_2^\alpha)) |\psi'(\theta c_1 + (1 - \theta)c_2)| d\theta \right] \\ & \leq \frac{c_2 - c_1}{2(c_2^\alpha - c_1^\alpha)} \left[\int_0^1 (((1 - \theta)c_1^{\alpha-1} + \theta c_2^{\alpha-1}) (\theta c_1 + (1 - \theta)c_2 - c_1^\alpha) |\psi'(\theta c_1 + (1 - \theta)c_2)| d\theta \right. \\ & \quad \left. + \int_0^1 (c_2^\alpha - ((1 - \theta)c_1^\alpha + \theta c_2^\alpha)) |\psi'(\theta c_1 + (1 - \theta)c_2)| d\theta \right] \\ & \leq \frac{c_2 - c_1}{2(c_2^\alpha - c_1^\alpha)} \left[\int_0^1 (((1 - \theta)c_1^{\alpha-1} + \theta c_2^{\alpha-1}) (\theta c_1 + (1 - \theta)c_2 - c_1^\alpha) \right. \\ & \quad \times [(1 - \theta)|\psi'(c_1)| + \theta|\psi'(c_2)|] d\theta \\ & \quad \left. + \int_0^1 (c_2^\alpha - ((1 - \theta)c_1^\alpha + \theta c_2^\alpha)) [(1 - \theta)|\psi'(c_1)| + \theta|\psi'(c_2)|] d\theta \right] \\ & = \frac{c_2 - c_1}{c_2^\alpha - c_1^\alpha} \left[\frac{1}{4} c_1^\alpha |\psi'(c_1)| + \frac{1}{12} c_1^\alpha |\psi'(c_2)| + \frac{1}{12} c_1 c_2^{\alpha-1} |\psi'(c_1)| + \frac{1}{12} c_1 c_2^{\alpha-1} |\psi'(c_2)| \right. \\ & \quad + \frac{1}{12} c_1^{\alpha-1} c_2 |\psi'(c_1)| + \frac{1}{12} c_1^{\alpha-1} c_2 |\psi'(c_2)| + \frac{1}{12} c_2^\alpha |\psi'(c_1)| + \frac{1}{4} c_2^\alpha |\psi'(c_2)| \\ & \quad - \frac{1}{2} c_1^\alpha |\psi'(c_1)| - \frac{1}{2} c_1^\alpha |\psi'(c_2)| + \frac{1}{2} c_2^\alpha |\psi'(c_1)| + \frac{1}{2} c_2^\alpha |\psi'(c_2)| - \frac{1}{3} c_1^\alpha |\psi'(c_1)| \\ & \quad \left. - \frac{1}{6} c_1^\alpha |\psi'(c_2)| - \frac{1}{6} c_2^\alpha |\psi'(c_1)| - \frac{1}{3} c_2^\alpha |\psi'(c_2)| \right] \\ & = \frac{c_2 - c_1}{2(c_2^\alpha - c_1^\alpha)} \left[\frac{(|\psi'(c_1)| + |\psi'(c_2)|)(5c_2^\alpha - 7c_1^\alpha + c_1c_2^{\alpha-1} + c_1^{\alpha-1}c_2)}{12} \right]. \quad \square \end{aligned}$$

Remark 2.4 Let $\alpha = 1$. Then inequality (2.2) becomes

$$\left| \frac{\psi(c_1) + \psi(c_2)}{2} - \frac{1}{c_2 - c_1} \int_{c_1}^{c_2} \psi(\xi) d\xi \right| \leq \frac{c_2 - c_1}{4} [|\psi'(c_1)| + |\psi'(c_2)|].$$

Theorem 2.5 Let $\alpha \in (0, 1]$, $q > 1$, $c_1, c_2 \in \mathbb{R}$ with $0 \leq c_1 < c_2$ and $\psi : [c_1, c_2] \rightarrow \mathbb{R}$ be an α -differentiable function on (c_1, c_2) . Then the inequality

$$\begin{aligned} & \left| \frac{\psi(c_1) + \psi(c_2)}{2} - \frac{\alpha}{c_2^\alpha - c_1^\alpha} \int_{c_1}^{c_2} \psi(\xi) d_\alpha \xi \right| \\ & \leq \frac{c_2 - c_1}{2(c_2^\alpha - c_1^\alpha)} \left[(A_1(\alpha))^{1-1/q} \{A_2(\alpha) |\psi'(c_1)|^q + A_3(\alpha) |\psi'(c_2)|^q\}^{1/q} \right. \\ & \quad \left. + (B_1(\alpha))^{1-1/q} \{B_2(\alpha) |\psi'(c_1)|^q + B_3(\alpha) |\psi'(c_2)|^q\}^{1/q} \right] \end{aligned} \quad (2.3)$$

is valid if $D_\alpha(\psi) \in L_\alpha^1([c_1, c_2])$ and $|\psi'|^q$ is convex on $[c_1, c_2]$, where

$$\begin{aligned} A_1(\alpha) &= \left[\frac{c_1^{\alpha+1} - c_2^{\alpha+1}}{(\alpha+1)(c_1 - c_2)} \right] - c_1^\alpha, & B_1(\alpha) &= c_2^\alpha - \left[\frac{c_1^{\alpha+1} - c_2^{\alpha+1}}{(\alpha+1)(c_1 - c_2)} \right], \\ A_2(\alpha) &= \left[\frac{-c_2^{\alpha+1}}{(\alpha+1)(c_1 - c_2)} + \frac{c_1^{\alpha+2} - c_2^{\alpha+2}}{(\alpha+1)(\alpha+2)(c_1 - c_2)^2} - \frac{c_1^\alpha}{2} \right], \\ B_2(\alpha) &= \left[\frac{c_2^\alpha}{2} + \frac{c_2^{\alpha+1}}{(\alpha+1)(c_1 - c_2)} + \frac{c_1^{\alpha+2} - c_2^{\alpha+2}}{(\alpha+1)(\alpha+2)(c_1 - c_2)^2} \right], \\ A_3(\alpha) &= \left[\frac{c_1^{\alpha+1}}{(\alpha+1)(c_1 - c_2)} - \frac{c_1^{\alpha+2} - c_2^{\alpha+2}}{(\alpha+1)(\alpha+2)(c_1 - c_2)^2} - \frac{c_1^\alpha}{2} \right], \\ B_3(\alpha) &= \left[\frac{c_2^\alpha}{2} - \frac{c_1^{\alpha+1}}{(\alpha+1)(c_1 - c_2)} + \frac{c_1^{\alpha+2} - c_2^{\alpha+2}}{(\alpha+1)(\alpha+2)(c_1 - c_2)^2} \right]. \end{aligned}$$

Proof From Lemma 2.1 and the well-known Hölder mean inequality together with the convexity of $|\psi'|^q$ on the interval $[c_1, c_2]$ we clearly see that

$$\begin{aligned} & \left| \frac{\psi(c_1) + \psi(c_2)}{2} - \frac{\alpha}{c_2^\alpha - c_1^\alpha} \int_{c_1}^{c_2} \psi(\xi) d_\alpha \xi \right| \\ &= \left| \frac{c_2 - c_1}{2(c_2^\alpha - c_1^\alpha)} \left[\int_0^1 ((\theta c_1 + (1-\theta)c_2)^{2\alpha-1} - c_1^\alpha (\theta c_1 + (1-\theta)c_2)^{\alpha-1}) \right. \right. \\ & \quad \times D_\alpha(\psi)(\theta c_1 + (1-\theta)c_2) d\theta \\ & \quad \left. \left. + \int_0^1 ((\theta c_1 + (1-\theta)c_2)^{2\alpha-1} - c_2^\alpha (\theta c_1 + (1-\theta)c_2)^{\alpha-1}) D_\alpha(\psi)(\theta c_1 + (1-\theta)c_2) d\theta \right] \right| \\ &\leq \frac{c_2 - c_1}{2(c_2^\alpha - c_1^\alpha)} \left[\int_0^1 ((\theta c_1 + (1-\theta)c_2)^\alpha - c_1^\alpha) |\psi'(\theta c_1 + (1-\theta)c_2)| d\theta \right. \\ & \quad \left. + \int_0^1 (c_2^\alpha - (\theta c_1 + (1-\theta)c_2)^\alpha) |\psi'(\theta c_1 + (1-\theta)c_2)| d\theta \right], \end{aligned} \quad (2.4)$$

$$\begin{aligned} & \int_0^1 ((\theta c_1 + (1-\theta)c_2)^\alpha - c_1^\alpha) |\psi'(\theta c_1 + (1-\theta)c_2)| d\theta \\ &\leq \left(\int_0^1 ((\theta c_1 + (1-\theta)c_2)^\alpha - c_1^\alpha) d\theta \right)^{1-1/q} \\ & \quad \times \left(\int_0^1 ((\theta c_1 + (1-\theta)c_2)^\alpha - c_1^\alpha) |\psi'(\theta c_1 + (1-\theta)c_2)|^q d\theta \right)^{1/q}, \end{aligned} \quad (2.5)$$

$$\begin{aligned} & \int_0^1 (c_2^\alpha - (\theta c_1 + (1-\theta)c_2)^\alpha) |\psi'(\theta c_1 + (1-\theta)c_2)| d\theta \\ & \leq \left(\int_0^1 (c_2^\alpha - (\theta c_1 + (1-\theta)c_2)^\alpha) d\theta \right)^{1-1/q} \\ & \quad \times \left(\int_0^1 (c_2^\alpha - (\theta c_1 + (1-\theta)c_2)^\alpha) |\psi'(\theta c_1 + (1-\theta)c_2)|^q d\theta \right)^{1/q}, \end{aligned} \quad (2.6)$$

$$\begin{aligned} & \int_0^1 ((\theta c_1 + (1-\theta)c_2)^\alpha - c_1^\alpha) |\psi'(\theta c_1 + (1-\theta)c_2)|^q d\theta \\ & \leq \int_0^1 ((\theta c_1 + (1-\theta)c_2)^\alpha - c_1^\alpha) [(1-\theta) |\psi'(c_1)|^q + \theta |\psi'(c_2)|^q] d\theta \\ & = |\psi'(c_1)|^q \int_0^1 ((\theta c_1 + (1-\theta)c_2)^\alpha - c_1^\alpha) (1-\theta) d\theta \\ & \quad + |\psi'(c_2)|^q \int_0^1 ((\theta c_1 + (1-\theta)c_2)^\alpha - c_1^\alpha) \theta d\theta \\ & = |\psi'(c_1)|^q \left[\frac{-c_2^{\alpha+1}}{(\alpha+1)(c_1-c_2)} + \frac{c_1^{\alpha+2} - c_2^{\alpha+2}}{(\alpha+1)(\alpha+2)(c_1-c_2)^2} - \frac{c_1^\alpha}{2} \right] \\ & \quad + |\psi'(c_2)|^q \left[\frac{\alpha+1}{(\alpha+1)(c_1-c_2)} - \frac{c_1^{\alpha+2} - c_2^{\alpha+2}}{(\alpha+1)(\alpha+2)(c_1-c_2)^2} - \frac{c_1^\alpha}{2} \right], \end{aligned} \quad (2.7)$$

$$\begin{aligned} & \int_0^1 (c_2^\alpha - (\theta c_1 + (1-\theta)c_2)^\alpha) |\psi'(\theta c_1 + (1-\theta)c_2)|^q d\theta \\ & \leq \int_0^1 (c_2^\alpha - (\theta c_1 + (1-\theta)c_2)^\alpha) [(1-\theta) |\psi'(c_1)|^q + \theta |\psi'(c_2)|^q] d\theta \\ & = |\psi'(c_1)|^q \int_0^1 (c_2^\alpha - (\theta c_1 + (1-\theta)c_2)^\alpha) (1-\theta) d\theta \\ & \quad + |\psi'(c_2)|^q \int_0^1 (c_2^\alpha - (\theta c_1 + (1-\theta)c_2)^\alpha) \theta d\theta \\ & = |\psi'(c_1)|^q \left[\frac{c_2^\alpha}{2} + \frac{c_2^{\alpha+1}}{(\alpha+1)(c_1-c_2)} + \frac{c_1^{\alpha+2} - c_2^{\alpha+2}}{(\alpha+1)(\alpha+2)(c_1-c_2)^2} \right] \\ & \quad + |\psi'(c_2)|^q \left[\frac{c_2^\alpha}{2} - \frac{c_1^{\alpha+1}}{(\alpha+1)(c_1-c_2)} + \frac{c_1^{\alpha+2} - c_2^{\alpha+2}}{(\alpha+1)(\alpha+2)(c_1-c_2)^2} \right]. \end{aligned} \quad (2.8)$$

Therefore, inequality (2.3) follows easily from (2.4)-(2.8). \square

Remark 2.6 Let $\alpha = 1$. Then inequality (2.3) becomes

$$\begin{aligned} & \left| \frac{\psi(c_1) + \psi(c_2)}{2} - \frac{1}{c_2 - c_1} \int_{c_1}^{c_2} \psi(\xi) d\xi \right| \\ & \leq \frac{1}{2} \left(\frac{c_2 - c_1}{2} \right)^{1-1/q} \left[\{A_2(1) |\psi'(c_1)|^q + A_3(1) |\psi'(c_2)|^q\}^{1/q} \right. \\ & \quad \left. + \{B_2(1) |\psi'(c_1)|^q + B_3(1) |\psi'(c_2)|^q\}^{1/q} \right] \end{aligned}$$

with

$$\begin{aligned} A_2(1) &= \frac{c_2 - c_1}{3}, & B_2(1) &= \frac{(c_1 + c_2)^2 + 2c_1c_2}{6(c_1 - c_2)}, \\ A_3(1) &= \frac{c_2 - c_1}{6}, & B_3(1) &= \frac{c_2 - c_1}{3}. \end{aligned}$$

Theorem 2.7 Let $\alpha \in (0, 1]$, $q > 1$, $c_1, c_2 \in \mathbb{R}$ with $0 \leq c_1 < c_2$ and $\psi : [c_1, c_2] \rightarrow \mathbb{R}$ be an α -differentiable function on (c_1, c_2) . Then the inequality

$$\left| \frac{\psi(c_1) + \psi(c_2)}{2} - \frac{\alpha}{c_2^\alpha - c_1^\alpha} \int_{c_1}^{c_2} \psi(\xi) d_\alpha \xi \right| \leq \frac{c_2 - c_1}{2(c_2^\alpha - c_1^\alpha)} \left[A_1(\alpha) \psi' \left(\frac{C_1(\alpha)}{A_1(\alpha)} \right) + B_1(\alpha) \psi' \left(\frac{C_2(\alpha)}{B_1(\alpha)} \right) \right] \quad (2.9)$$

holds if $D_\alpha(\psi) \in L_\alpha^1([c_1, c_2])$ and $|\psi'|^q$ is concave on $[c_1, c_2]$, where $A_1(\alpha)$ and $B_1(\alpha)$ are defined as in Theorem 2.5, and $C_1(\alpha)$ and $C_2(\alpha)$ are defined by

$$C_1(\alpha) = \left[\frac{c_1^{\alpha+2} - c_2^{\alpha+2}}{(\alpha+2)(c_1 - c_2)} - \frac{c_1^\alpha(c_1 - c_2)}{2} \right], \quad C_2(\alpha) = \left[\frac{c_2^\alpha(c_1 + c_2)}{2} - \frac{c_1^{\alpha+2} - c_2^{\alpha+2}}{(\alpha+2)(c_1 - c_2)} \right].$$

Proof It follows from the concavity of $|\psi'|^q$ and the Hölder mean inequality that

$$\begin{aligned} (\theta |\psi'(c_1)| + (1-\theta) |\psi'(c_2)|)^q &\leq \theta |\psi'(c_1)|^q + (1-\theta) |\psi'(c_2)|^q \leq |\psi'(\theta c_1 + (1-\theta)c_2)|^q, \\ |\psi'(\theta c_1 + (1-\theta)c_2)| &\geq \theta |\psi'(c_1)| + (1-\theta) |\psi'(c_2)|, \end{aligned}$$

which implies that $|\psi'|$ is also concave. Making use of Lemma 2.1 and the Jensen integral inequality, we have

$$\begin{aligned} &\left| \frac{\psi(c_1) + \psi(c_2)}{2} - \frac{\alpha}{c_2^\alpha - c_1^\alpha} \int_{c_1}^{c_2} \psi(\xi) d_\alpha \xi \right| \\ &= \left| \frac{c_2 - c_1}{2(c_2^\alpha - c_1^\alpha)} \left[\int_0^1 ((\theta c_1 + (1-\theta)c_2)^{2\alpha-1} - c_1^\alpha (\theta c_1 + (1-\theta)c_2)^{\alpha-1}) \right. \right. \\ &\quad \times D_\alpha(\psi)(\theta c_1 + (1-\theta)c_2) d\theta \\ &\quad \left. \left. + \int_0^1 ((\theta c_1 + (1-\theta)c_2)^{2\alpha-1} - c_2^\alpha (\theta c_1 + (1-\theta)c_2)^{\alpha-1}) D_\alpha(\psi)(\theta c_1 + (1-\theta)c_2) d\theta \right] \right| \\ &\leq \frac{c_2 - c_1}{2(c_2^\alpha - c_1^\alpha)} \left[\int_0^1 ((\theta c_1 + (1-\theta)c_2)^\alpha - c_1^\alpha) |\psi'(\theta c_1 + (1-\theta)c_2)| d\theta \right. \\ &\quad \left. + \int_0^1 (c_2^\alpha - (\theta c_1 + (1-\theta)c_2)^\alpha) |\psi'(\theta c_1 + (1-\theta)c_2)| d\theta \right], \quad (2.10) \end{aligned}$$

$$\begin{aligned} &\int_0^1 ((\theta c_1 + (1-\theta)c_2)^\alpha - c_1^\alpha) |\psi'(\theta c_1 + (1-\theta)c_2)| d\theta \\ &\leq \left(\int_0^1 ((\theta c_1 + (1-\theta)c_2)^\alpha - c_1^\alpha) d\theta \right) \\ &\quad \times \psi' \left(\frac{\int_0^1 ((\theta c_1 + (1-\theta)c_2)^\alpha - c_1^\alpha) (\theta c_1 + (1-\theta)c_2) d\theta}{\int_0^1 ((\theta c_1 + (1-\theta)c_2)^\alpha - c_1^\alpha) d\theta} \right) \\ &= A_1(\alpha) \psi' \left(\frac{C_1(\alpha)}{A_1(\alpha)} \right), \quad (2.11) \end{aligned}$$

$$\begin{aligned} &\int_0^1 (c_2^\alpha - (\theta c_1 + (1-\theta)c_2)^\alpha) |\psi'(\theta c_1 + (1-\theta)c_2)| d\theta \\ &\leq \left(c_2^\alpha - \int_0^1 ((\theta c_1 + (1-\theta)c_2)^\alpha) d\theta \right) \end{aligned}$$

$$\begin{aligned} & \times \psi' \left(\frac{\int_0^1 (c_2^\alpha - (\theta c_1 + (1-\theta)c_2)^\alpha) (\theta c_1 + (1-\theta)c_2) d\theta}{\int_0^1 (c_2^\alpha - (\theta c_1 + (1-\theta)c_2)^\alpha) d\theta} \right) \\ & = B_1(\alpha) \psi' \left(\frac{C_2(\alpha)}{B_1(\alpha)} \right). \end{aligned} \quad (2.12)$$

Therefore, inequality (2.9) follows easily from (2.10)-(2.12). \square

Remark 2.8 Let $\alpha = 1$. Then inequality (2.9) leads to

$$\begin{aligned} & \left| \frac{\psi(c_1) + \psi(c_2)}{2} - \frac{1}{c_2 - c_1} \int_{c_1}^{c_2} \psi(\xi) d\xi \right| \\ & \leq \frac{c_2 - c_1}{4} \left[\psi' \left(\frac{2c_2^2 - c_1^2 + 5c_1c_2}{3(c_2 - c_1)} \right) + \psi' \left(\frac{c_2^2 - 2c_1^2 + c_1c_2}{3(c_2 - c_1)} \right) \right]. \end{aligned}$$

3 Applications to special means of real numbers

Let $\alpha \in (0, 1]$, $r \in \mathbb{R}$, $r \neq 0, -\alpha$ and $a, b > 0$ with $a \neq b$. Then the arithmetic mean $A(a, b)$, logarithmic mean $L(a, b)$ and (α, r) th generalized logarithmic mean $L_{(\alpha, r)}(a, b)$ of a and b are defined by

$$A(a, b) = \frac{a+b}{2}, \quad L(a, b) = \frac{a-b}{\log a - \log b}, \quad L_{(\alpha, r)}(a, b) = \left[\frac{\alpha(b^{\alpha+r} - a^{\alpha+r})}{(\alpha+r)(b^\alpha - a^\alpha)} \right]^{1/r},$$

respectively. Then from Theorems 2.3 and 2.5 together with the convexities of the functions $\xi \rightarrow \xi^r$ and $\xi \rightarrow 1/\xi$ on the interval $(0, \infty)$ we get several new inequalities for the arithmetic, logarithmic and generalized logarithmic means as follows.

Theorem 3.1 Let $c_1, c_2 \in \mathbb{R}$ with $0 < c_1 < c_2$, $r > 1$, $q > 1$ and $\alpha \in (0, 1]$. Then we have

$$\begin{aligned} & |A(c_1^r, c_2^r) - L_{(\alpha, r)}^r(c_1, c_2)| \\ & \leq \frac{r(c_2 - c_1)(5c_2^\alpha - 7c_1^\alpha + c_1c_2^{\alpha-1} + c_1^{\alpha-1}c_2)}{12(c_2^\alpha - c_1^\alpha)} A(|c_1|^{r-1}, |c_2|^{r-1}), \\ & |A(c_1^r, c_2^r) - L_{(\alpha, r)}^r(c_1, c_2)| \\ & \leq \frac{r(c_2 - c_1)}{2(c_2^\alpha - c_1^\alpha)} \left[(A_1(\alpha))^{1-1/q} \{A_2(\alpha)|c_1|^{(r-1)q} + A_3(\alpha)|c_2|^{(r-1)q}\}^{1/q} \right. \\ & \quad \left. + (B_1(\alpha))^{1-1/q} \{B_2(\alpha)|c_1|^{(r-1)q} + B_3(\alpha)|c_2|^{(r-1)q}\}^{1/q} \right], \\ & |A(c_1^{-1}, c_2^{-1}) - L_{(\alpha, -1)}^{-1}(c_1, c_2)| \leq \frac{(c_2 - c_1)(5c_2^\alpha - 7c_1^\alpha + c_1c_2^{\alpha-1} + c_1^{\alpha-1}c_2)}{12(c_2^\alpha - c_1^\alpha)} A(c_1^{-2}, c_2^{-2}), \\ & |A(c_1^{-1}, c_2^{-1}) - L_{(\alpha, -1)}^{-1}(c_1, c_2)| \leq \frac{c_2 - c_1}{2(c_2^\alpha - c_1^\alpha)} \left[(A_1(\alpha))^{1-1/q} \{A_2(\alpha)|c_1|^{-2q} + A_3(\alpha)|c_2|^{-2q}\}^{1/q} \right. \\ & \quad \left. + (B_1(\alpha))^{1-1/q} \{B_2(\alpha)|c_1|^{-2q} + B_3(\alpha)|c_2|^{-2q}\}^{\frac{1}{q}} \right], \end{aligned}$$

where $A_1(\alpha)$, $A_2(\alpha)$, $A_3(\alpha)$, $B_1(\alpha)$, $B_2(\alpha)$ and $B_3(\alpha)$ are defined as in Theorem 2.5.

4 Applications to the trapezoidal formula

Let Δ be a division $c_1 = \xi_0 < \xi_1 < \cdots < \xi_{n-1} < \xi_n = c_2$ of the interval $[c_1, c_2]$ and consider the quadrature formula

$$\int_{c_1}^{c_2} \psi(\xi) d_\alpha \xi = T_\alpha(\psi, \Delta) + E_\alpha(\psi, \Delta),$$

where

$$T_\alpha(\psi, \Delta) = \sum_{i=0}^{n-1} \frac{\psi(\xi_i) + \psi(\xi_{i+1})}{2} \frac{(\xi_{i+1}^\alpha - \xi_i^\alpha)}{\alpha}$$

is the trapezoidal version and $E_\alpha(\psi, \Delta)$ denotes the associated approximation error. In this section, we are going to derive several new error estimations for the trapezoidal formula.

Theorem 4.1 *Let $\alpha \in (0, 1]$, $c_1, c_2 \in \mathbb{R}$ with $0 \leq c_1 < c_2$, $\psi : [c_1, c_2] \rightarrow \mathbb{R}$ be an α -differentiable function on (c_1, c_2) and Δ be a division $c_1 = \xi_0 < \xi_1 < \cdots < \xi_{n-1} < \xi_n = c_2$ of the interval $[c_1, c_2]$. Then the inequality*

$$|E_\alpha(\psi, \Delta)| \leq \frac{1}{12\alpha} \max\{|\psi'(c_1)|, |\psi'(c_2)|\} \sum_{i=0}^{n-1} (\xi_{i+1} - \xi_i) (5\xi_{i+1}^\alpha - 7\xi_i^\alpha + \xi_i \xi_{i+1}^{\alpha-1} + \xi_i^{\alpha-1} \xi_{i+1})$$

holds if $D_\alpha(\psi) \in L_\alpha^1([c_1, c_2])$ and $|\psi'|$ is convex on $[c_1, c_2]$.

Proof Applying Theorem 2.3 on the subinterval $[\xi_i, \xi_{i+1}]$ ($i = 0, 1, \dots, n-1$) of the division Δ , we have

$$\begin{aligned} & \left| \frac{\psi(\xi_i) + \psi(\xi_{i+1})}{2} \frac{(\xi_{i+1}^\alpha - \xi_i^\alpha)}{\alpha} - \int_{\xi_i}^{\xi_{i+1}} \psi(\xi) d_\alpha \xi \right| \\ & \leq \frac{(\xi_{i+1} - \xi_i)}{2\alpha} \left[\frac{(|\psi'(\xi_i)| + |\psi'(\xi_{i+1})|)(5\xi_{i+1}^\alpha - 7\xi_i^\alpha + \xi_i \xi_{i+1}^{\alpha-1} + \xi_i^{\alpha-1} \xi_{i+1})}{12} \right]. \end{aligned} \quad (4.1)$$

It follows from (4.1) and the convexity of $|\psi'(\xi)|$ on the interval $[c_1, c_2]$ that

$$\begin{aligned} |E_\alpha(\psi, \Delta)| &= \left| T_\alpha(\psi, \Delta) - \int_{c_1}^{c_2} \psi(\xi) d_\alpha \xi \right| \\ &= \left| \sum_{i=0}^{n-1} \left[\frac{\psi(\xi_i) + \psi(\xi_{i+1})}{2} \frac{(\xi_{i+1}^\alpha - \xi_i^\alpha)}{\alpha} - \int_{\xi_i}^{\xi_{i+1}} \psi(\xi) d_\alpha \xi \right] \right| \\ &\leq \sum_{i=0}^{n-1} \left| \frac{\psi(\xi_i) + \psi(\xi_{i+1})}{2} \frac{(\xi_{i+1}^\alpha - \xi_i^\alpha)}{\alpha} - \int_{\xi_i}^{\xi_{i+1}} \psi(\xi) d_\alpha \xi \right| \\ &\leq \frac{1}{2\alpha} \sum_{i=0}^{n-1} (\xi_{i+1} - \xi_i) \left[\frac{(|\psi'(\xi_i)| + |\psi'(\xi_{i+1})|)(5\xi_{i+1}^\alpha - 7\xi_i^\alpha + \xi_i \xi_{i+1}^{\alpha-1} + \xi_i^{\alpha-1} \xi_{i+1})}{12} \right] \\ &= \frac{1}{12\alpha} \sum_{i=0}^{n-1} (\xi_{i+1} - \xi_i) \left[\frac{(|\psi'(\xi_i)| + |\psi'(\xi_{i+1})|)(5\xi_{i+1}^\alpha - 7\xi_i^\alpha + \xi_i \xi_{i+1}^{\alpha-1} + \xi_i^{\alpha-1} \xi_{i+1})}{2} \right] \end{aligned}$$

$$\begin{aligned} &\leq \frac{1}{12\alpha} \sum_{i=0}^{n-1} (\xi_{i+1} - \xi_i) (5\xi_{i+1}^\alpha - 7\xi_i^\alpha + \xi_i \xi_{i+1}^{\alpha-1} + \xi_i^{\alpha-1} \xi_{i+1}) \max\{|\psi'(\xi_i)|, |\psi'(\xi_{i+1})|\} \\ &\leq \frac{1}{12\alpha} \max\{|\psi'(c_1)|, |\psi'(c_2)|\} \sum_{i=0}^{n-1} (\xi_{i+1} - \xi_i) (5\xi_{i+1}^\alpha - 7\xi_i^\alpha + \xi_i \xi_{i+1}^{\alpha-1} + \xi_i^{\alpha-1} \xi_{i+1}). \quad \square \end{aligned}$$

Making use of arguments analogous to the proof of Theorem 4.1, we get Theorem 4.2 immediately.

Theorem 4.2 *Let $\alpha \in (0, 1]$, $q > 1$, $c_1, c_2 \in \mathbb{R}$ with $0 \leq c_1 < c_2$, $\psi : [c_1, c_2] \rightarrow \mathbb{R}$ be an α -differentiable function on (c_1, c_2) and Δ be a division $c_1 = \xi_0 < \xi_1 < \dots < \xi_{n-1} < \xi_n = c_2$ of the interval $[c_1, c_2]$. Then the inequality*

$$\begin{aligned} |E_\alpha(\psi, \Delta)| &\leq \sum_{i=0}^{n-1} \frac{(\xi_{i+1} - \xi_i)}{2\alpha} \left[(A_1(\alpha))^{1-1/q} \{A_2(\alpha) |\psi'(\xi_i)|^q + A_3(\alpha) |\psi'(\xi_{i+1})|^q\}^{1/q} \right. \\ &\quad \left. + (B_1(\alpha))^{1-1/q} \{B_2(\alpha) |\psi'(\xi_i)|^q + B_3(\alpha) |\psi'(\xi_{i+1})|^q\}^{1/q} \right] \end{aligned}$$

holds if $D_\alpha(\psi) \in L_\alpha^1([c_1, c_2])$ and $|\psi'|^q$ is convex on $[c_1, c_2]$, where $A_1(\alpha)$, $A_2(\alpha)$, $A_3(\alpha)$, $B_1(\alpha)$, $B_2(\alpha)$ and $B_3(\alpha)$ are defined as in Theorem 2.5.

5 Conclusion

In this work, we find an identity and several Hermite-Hadamard type inequalities for conformable fractional integrals, present some new inequalities for the arithmetic, logarithmic and generalized logarithmic means of two positive real numbers and provide the error estimations for the trapezoidal formula.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

All authors contributed equally to the writing of this paper. All authors read and approved the final manuscript.

Author details

¹Department of Mathematics, Huzhou University, Huzhou, 313000, China. ²Department of Mathematics, University of Peshawar, Peshawar, 25000, Pakistan. ³College of Engineering and Science, Victoria University, Melbourne, VIC 8001, Australia.

Acknowledgements

The research was supported by the Natural Science Foundation of China (Grants Nos. 61673169, 61374086, 11371125, 11401191) and the Tianyuan Special Funds of the National Natural Science Foundation of China (Grant No. 11626101).

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Received: 8 October 2016 Accepted: 20 April 2017 Published online: 28 April 2017

References

- Hermite, C: Sur deux limites d'une intégrale définie. *Mathesis* **3**, 82 (1883)
- Hadamard, J: Étude sur les propriétés des fonctions entières et en particulier d'une fonction considérée par Riemann. *J. Math. Pures Appl.* **58**, 171-215 (1893)
- Niculescu, CP, Persson, L-E: *Convex Functions and Their Applications*. Springer, New York (2006)
- Wang, M-K, Li, Y-M, Chu, Y-M: Inequalities and infinite product formula for Ramanujan generalized modular equation function. *Ramanujan J.* (2017). doi:10.1007/s11139-017-9888-3
- Wang, M-K, Chu, Y-M: Refinements of transformation inequalities for zero-balanced hypergeometric functions. *Acta Math. Sci.* **37B**(3), 607-622 (2017)

6. Adil Khan, M, Khurshid, Y, Ali, T: Hermite-Hadamard inequality for fractional integrals via η -convex functions. *Acta Math. Univ. Comen.* **86**(1), 153-164 (2017)
7. Adil Khan, M, Khurshid, Y, Ali, T, Rehman, N: Inequalities for three times differentiable functions. *Punjab Univ. J. Math.* **48**(2), 35-48 (2016)
8. Chu, Y-M, Adil Khan, M, Khan, TU, Ali, T: Generalizations of Hermite-Hadamard type inequalities for MT-convex functions. *J. Nonlinear Sci. Appl.* **9**(6), 4305-4316 (2016)
9. Wu, Y, Qi, F, Niu, D-W: Integral inequalities of Hermite-Hadamard type for the product of strongly logarithmically convex and other convex functions. *Maejo Int. J. Sci. Technol.* **9**(3), 394-402 (2015)
10. Noor, MA, Noor, KI, Awan, MU: Hermite-Hadamard inequalities for relative semi-convex functions and applications. *Filomat* **28**(2), 221-230 (2014)
11. Bai, R-F, Qi, F, Xi, B-Y: Hermite-Hadamard type inequalities for the m - and (α, m) -logarithmically convex functions. *Filomat* **27**(1), 1-7 (2013)
12. Matloka, M: On some Hadamard-type inequalities for (h_1, h_2) -preinvex functions on the co-ordinates. *J. Inequal. Appl.* **2013**, Article ID 227 (2013)
13. Zhang, X-M, Chu, Y-M, Zhang, X-H: The Hermite-Hadamard type inequality of GA-convexity functions. *J. Inequal. Appl.* **2010**, Article ID 507560 (2010)
14. Chu, Y-M, Wang, G-D, Zhang, X-H: Schur convexity and Hadamard's inequality. *Math. Inequal. Appl.* **13**(4), 725-731 (2010)
15. Bombardelli, M, Varošanec, S: Properties of h -convex functions related to the Hermite-Hadamard-Fejér inequalities. *Comput. Math. Appl.* **58**(9), 1869-1877 (2009)
16. Sarikaya, MZ, Saglam, A, Yildirim, H: On some Hadamard-type inequalities for h -convex functions. *J. Math. Inequal.* **2**(3), 335-341 (2008)
17. Kirmaci, US, Klaričić Bakula, M, Özdemir, ME, Pečarić, J: Hadamard-type inequalities for s -convex functions. *Appl. Math. Comput.* **193**(1), 26-35 (2007)
18. Noor, MA: On Hadamard integral inequalities involving two log-preinvex functions. *JIPAM. J. Inequal. Pure Appl. Math.* **8**(3), Article ID 75 (2007)
19. Dragomir, SS, McAndrew, A: Refinements of the Hermite-Hadamard inequality for convex functions. *JIPAM. J. Inequal. Pure Appl. Math.* **6**(5), Article ID 140 (2005)
20. Dragomir, SS, Fitzpatrick, S: The Hadamard inequalities for s -convex functions in the second sense. *Demonstr. Math.* **32**(4), 687-696 (1999)
21. Dragomir, SS, Pečarić, JE, Persson, L-E: Some inequalities of Hadamard type. *Soochow J. Math.* **21**(5), 335-341 (1995)
22. Dragomir, SS: Two mappings in connection to Hadamard's inequalities. *J. Math. Anal. Appl.* **167**(1), 49-56 (1992)
23. Dragomir, SS, Agarwal, RP: Two inequalities for differentiable mappings and applications to special means of real numbers and to trapezoidal formula. *Appl. Math. Lett.* **11**(5), 91-95 (1998)
24. Pearce, CEM, Pečarić, JE: Inequalities for differentiable mapping with application to special means and quadrature formulae. *Appl. Math. Lett.* **13**(2), 51-55 (2000)
25. Iyiola, OS, Nwaeze, ER: Some new results on the new conformable fractional calculus with application using D'Alambert approach. *Prog. Fract. Differ. Appl.* **2**(2), 115-122 (2016)
26. Abdeljawad, T: On conformable fractional calculus. *J. Comput. Appl. Math.* **279**, 57-66 (2015)
27. Hammad, MA, Khalil, R: Conformable fractional heat differential equations. *Int. J. Pure Appl. Math.* **94**(2), 215-221 (2014)
28. Hammad, MA, Khalil, R: Abel's formula and Wronskian for conformable fractional differential equations. *Int. J. Differ. Equ. Appl.* **13**(3), 177-183 (2014)
29. Khalil, R, Al Horani, M, Yousef, A, Sababheh, M: A new definition of fractional derivative. *J. Comput. Appl. Math.* **264**, 65-70 (2014)
30. Cheng, J-F, Chu, Y-M: On the fractional difference equations of order $(2, q)$. *Abstr. Appl. Anal.* **2011**, Article ID 497259 (2011)
31. Cheng, J-F, Chu, Y-M: Solution to the linear fractional differential equation using Adomian decomposition method. *Math. Probl. Eng.* **2011**, Article ID 587068 (2011)
32. Cheng, J-F, Chu, Y-M: Fractional difference equations with real variable. *Abstr. Appl. Anal.* **2012**, Article ID 918529 (2012)
33. Anderson, DR: Taylor's formula and integral inequalities for conformable fractional derivatives. In: *Contributions in Mathematics and Engineering, in Honor of Constantin Carathéodory*. Springer, Berlin (2016)

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Immediate publication on acceptance
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► springeropen.com